

A bioarchaeological approach to the Iron Age in Switzerland: stable isotope analyses ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$) of human remains

Negahnaz Moghaddam¹ · Felix Müller² · Sandra Lösch¹

Received: 3 August 2016 / Accepted: 17 November 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract In Switzerland, a large number of Iron Age burial sites were found in the last century. Changes in living conditions and socio-cultural behavior may have occurred over time and space and could be reflected in the dietary habits, social stratigraphy within populations and migration patterns. This study attempts to shed light on these aspects with the application of stable isotope analyses. Human remains from 11 different burial sites ($n = 164$) in the area of today's Swiss Plateau and Swiss Alpine regions were investigated. Temporal and geographical variations as well as sex and age-related dietary differences were analyzed through isotopic studies ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$). In total, the data of 129 individuals could be evaluated. Highly significant differences between the burial sites were found, with higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the Alpine regions. Cultural and/or climatic changes as well as the different geological conditions might have led to distinct patterns of crop cultivation and animal husbandry and consequently to significantly different dietary habits in the Plateau and the Alpine regions. The data indicate a higher intake of millet and animal protein including early dairy production in the southern regions, probably influenced by the Mediterranean world.

Electronic supplementary material The online version of this article (doi:10.1007/s12520-016-0441-x) contains supplementary material, which is available to authorized users.

✉ Sandra Lösch
Sandra.Loesch@irm.unibe.ch

¹ Department of Physical Anthropology, Institute of Forensic Medicine, University of Bern, Sulgenauweg 40, 3007 Bern, Switzerland

² Department of Prehistory, Institute of Archaeological Sciences, University of Bern, Bern, Switzerland

Cultural exchange between geographical regions might have been facilitated by migration during the Iron Age as suggested by the $\delta^{34}\text{S}$.

Keywords Stable isotopes · Paleodiet · Iron Age

Introduction

Bioarchaeological approaches to Late Iron Age human remains have become more and more important for anthropological and archaeological research (Knipper et al. 2014; Koon and Nicholls 2016; Le Huray and Schutkowski 2005; Lightfoot et al. 2015; Moghaddam et al. 2016; Naumann et al. 2014; Oelze et al. 2012; Redfern et al. 2010). Due to the poor preservation of many Iron Age skeletons and the lack of written sources from that time, stable isotope analyses provide important new insights into the Iron Age dietary and migration patterns (Le Huray et al. 2006).

The Iron Age in central Europe is characterized by the Hallstatt (Early Iron Age from 800 to 450 BC) and the La Tène period (Late Iron Age from 450 to 15 BC). The La Tène period is subdivided into the stages LT A (Early La Tène) to LT D (Late La Tène). The foundation of the City of Massalia around 600 BC, nowadays known as Marseille, played a major role in the history of the “Celts” (Müller et al. 1999; Müller and Lüscher 2004). It influenced the Iron Age culture including the greater tribes near the Rhône such as the Sedunii in the area of today's Valais in the south of Switzerland (Curdy et al. 2009). Between the fourth and third centuries BC, La Tène culture expanded in most areas of central Europe including migration waves to Italy, the Balkans and to today's Turkey (Kaenel 1999). Iron Age populations were a diverse group of small-scale societies with similar religion, language, and culture. The term “Celts” should

therefore be used carefully. It remains unclear whether this term defines populations with the same language or also political groups (Collis 1984). One of the greater tribes was the Helvetii in the Swiss Plateau. Other tribes in the area of modern Switzerland were the Allobroges around Geneva, Raurici and Latobrigi in modern Basel and Schaffhausen, the Leponti in Ticino, and the Raeti in Grisons (Müller et al. 1999). “Celtic” culture, however, rapidly declined after the Roman Empire conquered Spain, Gaul, and areas of the upper Danube.

In the area of Switzerland, a large amount of Iron Age burial sites were excavated in the last century and have raised questions: how was the social structure within populations organized? Were there any differences in the socioeconomic and cultural structure between societies and within a population?

This study presents stable isotope data ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$) of human remains from different regions in Switzerland in order to obtain more information about “Celtic” populations. It is suggested that changes and differences in living conditions and socio-cultural behavior might have taken place through time and space. This might be reflected in the dietary habits, social stratigraphy within populations and possible migration of individuals. Therefore, this study aimed to analyze geographical and temporal dietary differences of individuals from Late Iron Age burial sites in Switzerland. Burials found in the Swiss Plateau and the Swiss Alpine regions were compared to discover whether different geological conditions and climate might have influenced living conditions and economical structures during the period. Cultural changes during the Swiss Late Iron Age are evident through the archaeological findings. Archaeological research shows differences in grave goods and burial rites between the Swiss Plateau and the southern area of Switzerland. Although no items of weaponry were found from the LT C period onwards in a larger necropolis in the area of Bern, weapons as grave goods were found in the burial sites of Sion dating to the LT D period (Jud and Ulrich-Bochsler 2014). In addition, dietary differences should now be brought to light through bioarchaeological studies.

Stable isotope analyses: the reconstruction of diet and migration

The technique of stable isotope ratio analysis is a frequently used tool to assess dietary habits, to obtain information about social stratification and mobility of past populations (Dupras and Schwarcz 2001; Fuller et al. 2012; Jay et al. 2013; Jay and Richards 2006; Katzenberg 2008; Knipper et al. 2016; Lösch et al. 2006).

Stable carbon and nitrogen isotope analyses are the most common methods to examine diet (Katzenberg et al. 2012; Schoeninger and DeNiro 1984; Stevens and Hedges 2004). Analysis of $\delta^{13}\text{C}$ provides information about the different

plant sources in the diet: C_3 and C_4 plants differ in their photosynthetic pathway and therefore show different $\delta^{13}\text{C}$ values (Ambrose 1993; Ambrose and Norr 1993; Hoefs 2009). Due to a strong discrimination against ^{13}C the $\delta^{13}\text{C}$ values in C_3 plants such as wheat and barley are more negative (−19 to −35‰) than in C_4 plants (−9 to −35‰) (Ambrose 1993; Lee-Thorp 2008). The $\delta^{13}\text{C}$ value becomes enriched by +1‰ for each trophic level, from prey to predator collagen (Redfern et al. 2012).

Additionally, analysis of $\delta^{15}\text{N}$ can be used to predict a relative animal protein intake. Towards the top of the food chain, the $\delta^{15}\text{N}$ increases (“trophic level effect”) and therefore the variation of stable nitrogen isotope ratios provide information about trophic level (Richards et al. 1998). In bone collagen, the approximate shift from one trophic level to the next one is about +3 to +5‰ (Bocherens and Drucker 2003; Hedges and Reynard 2007). Individuals with mostly animal protein in their diet have higher $\delta^{15}\text{N}$ values than those consuming mainly plants (Richards et al. 1998; Stevens et al. 2010). Furthermore, $\delta^{15}\text{N}$ values show the relative contribution of terrestrial and marine resources (Schoeninger and DeNiro 1984).


The analysis of $\delta^{34}\text{S}$ has become important for stable isotope research for analyses of dietary habits and migration (Bollongino et al. 2013; Craig et al. 2006; Fornander et al. 2008; Kinaston et al. 2013; Moghaddam et al. 2016; Nehlich et al. 2011, 2012; Oelze et al. 2012; Privat et al. 2007; Richards et al. 2001, 2003; Vika 2009). In bone collagen, the $\delta^{34}\text{S}$ is passed along the food chain with just a small fractionation of approximately −1‰. Organisms in marine ecosystems have stable sulfur isotope ratios around +20‰ while terrestrial mammals have values even lower than +10‰ (Nehlich et al. 2012; Richards et al. 2003). However, the values of freshwater and terrestrial ecosystems range from −22‰ to +22‰ (Oelze et al. 2012; Privat et al. 2007). Additionally, the $\delta^{34}\text{S}$ values vary between geographical regions due to different geological conditions (Vika 2009). Therefore, sulfur isotopic signatures are influenced by dietary habits and location (Howcroft et al. 2012; Nehlich et al. 2011).

Material

In total, samples of 164 human and 5 animal bones were collected from 11 sites: 7 sites located on the Swiss Plateau and 4 sites in the Swiss Alps (Fig. 1; Table 1).

The animal bones from Reichenbachstrasse were identified as one pig, one cattle and two samples from ovicaprids (Rehazek and Nussbaumer 2014). One ovicaprid sample from Bonaduz in the area of Grisons was included (Table 2).

Fig. 1 Switzerland (provided by the Archaeological Service Bern; Source: Swisstopo): Major landscapes shown as *I* Jura Mountains, *II* Swiss Plateau, *III* Swiss Alps. Burial sites are indicated with numbers (*1* Engehalbinsel, *2* Ipsach, *3* Belp, *4* Niederwichttrach, *5* Bümpliz, *6* Stettlen-Deisswil, *7* Andelfingen, *8* Sion, *9* Bramois, *10* Castaneda, *11* Trun Darvella). Additionally, burial site of Münsingen is marked by *12*



Iron Age burial sites at the Swiss plateau

Today, the Swiss Plateau between the Jura Mountains and the Alps covers about 30% of Switzerland and elongates from Geneva in the south west to the German border in the north east (Fig. 1). Even though the area is flat in some parts, it shows mainly hilly areas, several large lakes (such as Lakes Geneva, Neuchâtel and Zurich) and rivers. The city of Bern

and the surrounding area provide the largest amount of skeletal material of Late Iron Age Switzerland. During the Iron Age, the Aare valley between the city of Bern and Lake Thun seems to have been part of a fairly well-developed settlement area that reached out to the West (Müller 1996). The Enge peninsula (Engehalbinsel) held the Celtic oppidum “Brenodor”, a city-like settlement, between the third and first centuries BC (Jud and Ulrich-Bochsler 2014). For some burial

Table 1 Summary of all burial sites: burial sites are indicated by numbers (*1* Engehalbinsel, *2* Ipsach, *3* Belp, *4* Niederwichttrach, *5* Bümpliz, *6* Stettlen-Deisswil, *7* Andelfingen, *8* Sion, *9* Bramois, *10* Castaneda, *11* Trun Darvella)

	Region	Burial site	Number of samples	Archaeological dating	Primary sources and additional literature
Swiss Plateau	Bern	1: Engehalbinsel	38	LT C=D	Hug (1956); Jud and Ulrich-Bochsler (2014)
		2: Ipsach	5	LT B–C	Ramstein (2010); Zweifel (2015)
		3: Belp	7	LT C–D	Schoch and Ulrich-Bochsler (1987); Suter and Ulrich-Bochsler (1984)
		4: Niederwichttrach	5	LT B–C	Schoch and Ulrich-Bochsler (1987); Stöckli (1995)
		5: Bümpliz	11	LT B–C	Hug (1956); Schoch and Ulrich-Bochsler (1987); Stähli (1977)
		6: Stettlen - Deisswil	8	LT B–C	Rey (1999)
	Zurich	7: Andelfingen	16	LT B–C	Viollier (1912)
Swiss Alps	Valais	Total	90	LT B–D	
		8: Sion	53	LT B–D	Curdy et al. (2009); Debard (2014)
		9: Bramois	17	LT B–D	Curdy et al. (2009); Debard (2014)
	Grisons	10: Castaneda	1	LT (A)B–C?	Keller-Tarnuzzer (1933); Nagy (2008)
		11: Trun Darvella	3	LT B–D	Tanner (1980)
		Total	74	LT B–D	

Table 2 Anthropological and stable isotope data of human ($n = 164$) and faunal ($n = 5$) remains; excluded samples are shown in italics and classification of dating group through ^{14}C analyses in bold.

Burial sites	Number	Age	Age class	Sex	Sample	Archaeological dating	^{14}C dating (2)	Grave goods	$\delta^{13}\text{C}$ [‰]V-PDB	$\delta^{15}\text{N}$ [‰]AIR	$\delta^{34}\text{S}$ [‰]V-CDT	%coll	%C	%N	%S	C/ N mol	C/S	N/S
Bern: Engenhalsinsel (Reichenbachstrasse)	A5182	46–51	Mature	f	Skull	D1			–18.9	7.8	5.1	2.6	40.7	15.5	0.1	3.1	313.1	119.2
	A5184	8–10	Infant II	nd	Long bone	/			–19.5	9.1	6.7	4.8	37.0	17.4	0.1	2.5	370.0	174.0
	A5185/13	8–9	Infant II	nd	Skull	/			–19.6	6.3	6.7	7.3	39.0	16.8	0.1	2.7	354.5	152.7
	A5185/16	23–40	Adult	f	Skull	C2			–19.8	6.9	6.8	5.6	41.8	16.5	0.1	3.0	380.0	150.0
	A5187	7–9	Infant II	nd	Skull	/			–20.2	7.8	5.6	1.0	36.6	12.8	0.1	3.3	406.7	142.2
	A5188	10–11	Infant II	nd	Skull	C2		Gold or bronze ring and/or pearl	–20.0	6.2	6.9	4.4	38.8	16.3	0.1	2.8	352.7	148.2
	A5189	30–50	Mature	f	Skull	C2			–19.6	8.1	6.6	3.9	41.9	16.8	0.1	2.9	299.3	120.0
	A5190	2–3	Infant	nd	Skull	C2		Gold or bronze ring and/or pearl	–19.8	8.3	6.6	7.1	38.6	16.7	0.1	2.7	350.9	151.8
	A5191	18–30	Adult	f	Skull	C2		Gold or bronze ring and/or pearl	–20.1	6.6	7.2	4.1	36.9	16.0	0.1	2.7	335.5	145.5
	A5192	30–40	Adult	f	Skull	D1		Gold or bronze ring and/or pearl	–19.1	8.0	4.7	4.0	45.0	16.8	0.1	3.1	321.4	120.0
	A5193	3–4	Infant	nd	Long bone	/			–19.7	6.6	6.8	3.8	39.8	16.8	0.1	2.8	306.2	129.2
	A5194	4–5	Infant	nd	Skull	D1			–19.9	7.5	6.0	4.7	40.5	16.9	0.1	2.8	337.5	140.8
	A5195	4–5	Infant	nd	Skull	/			–20.2	8.2	5.4	2.7	41.6	15.6	0.1	3.1	297.1	111.4
	A5196	8–11	Infant II	nd	Long bone	C/D			–20.1	7.0	5.8	2.3	42.3	17.1	0.1	2.9	352.5	142.5
	A5197	37–33	Adult	f	Skull	C2			–19.7	7.1	6.7	4.7	44.0	16.9	0.1	3.0	366.7	140.8
	A5198	4–5	Infant	nd	Skull	C2			–19.8	7.6	6.3	6.0	42.6	16.2	0.1	3.1	327.7	124.6
	A5199	37–46	Adult	f	Skull	C2		Gold or bronze ring and/or pearl	–19.6	8.1	6.0	4.5	39.9	15.4	0.1	3.0	306.9	118.5
	A5200	30–40	Adult	nd	Skull	C2		Coins	/	/	/	/	/	/	/	/	/	/
	A5201	54–60	Mature	f	Skull	/			–18.8	4.8	7.1	1.5	32.7	16.1	0.1	2.4	327.0	161.0
	A5202	23–30	Adult	f	Skull	D1		Gold or bronze ring and/or pearl	–19.2	7.6	5.9	1.7	41.9	15.1	0.1	3.2	419.0	151.0
	A5203	4–5	Infant	nd	Skull	D1			–19.5	8.0	6.4	5.2	41.3	16.0	0.1	3.0	375.5	145.5
	A5204	3–5	Infant	nd	Skull	D1			–19.5	8.4	6.3	5.7	42.0	16.4	0.1	3.0	323.1	126.2
	A5205	3–4	Infant	nd	Skull	C/D			–19.7	8.6	6.3	5.8	43.1	17.1	0.1	2.9	359.2	142.5
	A5206	23–40	Adult	f	Skull	/			–19.8	8.0	6.6	6.0	41.4	16.2	0.1	3.0	345.0	135.0
	A5207	23–40	Adult	f	Skull	C2		Gold or bronze ring/pearl	–19.5	6.9	7.1	4.1	43.2	17.3	0.1	2.9	360.0	144.2
	A5208	30+	Adult	f	Skull	C2		Coins and gold or bronze ring and/or pearl	/	/	/	/	/	/	/	/	/	/
	A5209	40+	Mature	f	Skull	C2		Coins	–19.3	8.8	5.4	4.2	42.1	15.8	0.1	3.1	323.8	121.5
	A5210	0.5–1	Infant	nd	Skull	D1			–18.7	11.3	6.7	3.3	39.7	14.6	0.1	3.2	397.0	146.0
	A5211	3–4	Infant	nd	Skull	D1			–19.3	7.1	7.0	5.7	33.0	16.1	0.1	2.4	330.0	161.0
	A5212	30–50	Adult	nd	Skull	D1		Coins	–19.3	7.0	7.0	4.2	39.4	16.6	0.1	2.8	328.3	138.3
	A5213	4–5	Infant	nd	Skull	/			–19.5	10.4	5.3	2.4	41.1	15.4	0.1	3.1	342.5	128.3
	A5214	0.5–1	Infant	nd	Skull	C2			–19.1	10.3	6.6	4.5	41.8	15.3	0.1	3.2	418.0	153.0
	A16	30–50	Adult	f	Femur	/			–18.2	8.1	5.4	5.1	44.3	16.4	0.1	3.2	316.4	117.1

Table 2 (continued)

Burial sites	Number	Age	Age class	Sex	Sample	Archaeological dating	¹⁴ C dating (2)	Grave goods	δ ¹³ C [‰]V-PDB	δ ¹⁵ N [‰]AIR	δ ³⁴ S [‰]V-CDT	%coll	%C	%N	%S	C/ N mol	C/S	N/S
Bern: Engehalbinsel (Rossfeldstrasse)	A17	20–30	Adult	m	Skull	/	/		–19.8	7.4	3.3	2.0	42.8	16.3	0.1	3.1	305.7	116.4
	A18	20–30	Adult	m	Ulna	/	/		–19.0	7.5	5.6	3.9	42.4	16.1	0.1	3.1	302.9	115.0
	A19	20–40	Adult	f	Skull	/	/		–19.7	8.2	1.6	1.4	41.7	16.1	0.1	3.0	320.8	123.8
	A20	30–40	Adult	m	Skull	/	/		–18.7	7.2	6.0	1.5	33.7	15.2	0.1	2.6	306.4	138.2
	A21	21–25	Adult	f	Ulna	/	/		–18.7	7.8	5.9	4.3	44.1	16.5	0.1	3.1	315.0	117.9
Bern: Ipsach	IP14	35–45	Adult	f	Skull	C1	/		–18.0	8.9	6.7	3.5	34.8	12.1	0.1	3.4	316.4	110.0
	IP50	14–20	Juvenile	f	Skull	B	/		/	/	/	/	/	/	/	/	/	/
	IP73	4–6	Infant	nd	Skull	B2	/		–19.2	10.7	6.2	3.2	26.5	9.0	0.1	3.4	240.9	81.8
	IP133	2.5–3.5	Infant	nd	Skull	B1/B2	/		–16.3	10.2	3.6	2.3	38.3	13.5	0.1	3.3	319.2	112.5
	IP136	5–7	Infant	nd	Skull	B2	/		–18.1	9.7	6.2	2.7	32.2	11.1	0.1	3.4	292.7	100.9
Bern: Belp	BEB-A4	20–30	Adult	m	Skull	C	196–55		–20.0	8.3	4.0	3.9	45.1	16.1	0.2	3.3	205.0	73.2
	BEB-A2268	45–60	Mature	f	Skull	C-D	BC		–18.8	9.3	1.8	2.1	43.3	15.0	0.2	3.4	206.2	71.4
	BEB-A2269	40+	Mature	m	Long bone	C-D	201–55		–19.1	7.8	0.2	1.6	44.9	16.3	0.2	3.2	224.5	81.5
	BEB-A2270	45–60	Mature	m	Skull	C	BC		–18.5	8.1	3.3	3.6	44.4	15.9	0.2	3.3	233.7	83.7
	BEB-A2271	40–55	Mature	m	Skull	C-D	348–94		–19.6	8.6	5.1	1.5	44.0	15.9	0.2	3.2	209.5	75.7
Bern: Niederwichterach	BEB-A2272	25–35	Adult	f	Skull	C	BC		–16.1	8.3	3.7	1.9	42.6	15.4	0.2	3.2	224.2	81.1
	BEB-A2276	30–50	Adult	f	Maxilla	/	/		–18.9	8.7	1.9	3.8	41.1	14.6	0.2	3.3	216.3	76.8
	NW-A1191	45–60	Mature	m	Skull	/	/		–20.6	9.0	3.9	1.9	29.3	10.2	0.2	3.3	172.4	60.0
	NW-A1189	25–40	Adult	f	Skull	B-C (D)	363–203		–19.7	8.8	2.0	1.9	42.9	15.0	0.2	3.3	204.3	71.4
	NW-A1190	20–40	Adult	nd	Mandibula	/	BC		–21.5	8.8	2.3	3.3	43.5	14.3	0.2	3.5	241.7	79.4
Bern: Bümpliz	NW-A1188	55–70	Senile	nd	Skull	B-C	374–204		–20.1	8.8	2.5	4.1	44.0	15.0	0.2	3.4	220.0	75.0
	NW-A2415	1–3	Infant	nd	Skull	/	BC		–19.8	8.8	1.3	2.3	35.9	12.1	0.2	3.5	188.9	63.7
	BBÜ-A6	25–40	Adult	f	Skull	A	385–206		–19.0	7.9	6.4	2.5	40.8	14.9	0.2	3.2	226.7	82.8
	BBÜ-A7	20–30	Adult	f	Skull	B-C	BC		–18.4	7.3	6.5	4.8	44.2	16.0	0.2	3.2	294.7	106.7
	BBÜ-A8	30–50	Adult	f	Skull	/	BC		–18.2	7.7	6.4	4.3	44.4	16.0	0.1	3.2	317.1	114.3
Bern: Bümpliz	BBÜ-A9	30–45	Adult	f	Skull	B-C	389–209		–18.0	7.3	6.7	4.3	43.9	15.8	0.1	3.2	337.7	121.5
	BBÜ-A10	55–70	Mature	f	Skull	/	BC		–19.7	8.4	2.1	2.6	44.8	16.3	0.1	3.2	373.3	135.8
	BBÜ-A11	20–40	Adult	f	Skull	C2	/		–17.7	7.0	6.7	2.4	43.6	15.7	0.1	3.2	311.4	112.1
	BBÜ-A12	20–40	Adult	m	Skull	B-C	388–208		–17.8	7.4	7.1	2.0	44.0	16.1	0.1	3.2	314.3	115.0
	BBÜ-A13	20–35	Adult	f	Skull	B1	BC		–19.1	8.4	5.4	3.0	42.2	15.3	0.1	3.2	301.4	109.3
Bern: Stettlen-Deisswil	BBÜ-A375	30–40	Adult	f	Skull	B-C	360–176		–17.9	7.6	6.8	5.2	44.5	16.2	0.2	3.2	296.7	108.0
	BBÜ-A376	14–18	Juvenile	nd	Skull	B-C	BC		–17.6	7.2	6.9	3.2	45.2	16.3	0.1	3.2	347.7	125.4
	BBÜ-A833	3–4	Infant	nd	Skull	A/B	382–204		–20.3	8.2	6.2	2.5	43.9	15.5	0.2	3.3	199.5	70.5
	SD-A126	25–40	Adult	f	Skull	C	/		–19.6	8.9	1.9	7.1	45.2	16.5	0.2	3.2	265.9	97.1

Table 2 (continued)

Burial sites	Number	Age	Age class	Sex	Sample	Archaeological dating	¹⁴ C dating (2)	Grave goods	δ ¹³ C [‰]V-PDB	δ ¹⁵ N [‰]AIR	δ ³⁴ S [‰]V-CDT	%coll	%C	%N	%S	C/N	C/S	N/S
Zürich: Andelfingen	SD-A127	40–60	Mature	f	Skull	C	365–198 BC		–18.6	8.5	1.4	5.6	44.1	15.9	0.2	3.2	294.0	106.0
	SD-A128	40–50	Mature	f	Skull	B-C			–19.0	8.6	2.6	1.4	40.3	14.3	0.2	3.3	212.1	75.3
	SD-A129	40–55	Mature	f	Skull	B		Gold or bronze ring and/or pearl	–19.9	10.4	1.3	5.0	43.8	15.9	0.1	3.2	312.9	113.6
	SD-A775	6–7	Infant lb	id	Skull	C			–18.7	8.6	2.1	7.0	43.2	15.7	0.1	3.2	308.6	112.1
	SD-A776	14–18	Juvenile	nd	Skull	B-C	365–203 BC		–18.6	8.2	3.1	6.3	46.2	16.9	0.2	3.2	243.2	88.9
	SD-A130	40–55	Mature	m	Tibia	B		Weapon	–19.9	9.1	2.2	5.7	45.2	16.3	0.2	3.2	226.0	81.5
	SD-A131	40–60	Mature	m	Skull	C			–19.9	9.0	3.8	5.7	44.9	15.8	0.2	3.3	299.3	105.3
	ZAF-4	25+	Adult	m	Femur	C			–18.1	8.8	8.5	1.9	43	15.3	0.1	3.3	390.9	139.1
	ZAF-5	25–40	Adult	f	Femur	C			–17.1	8.4	8.4	1.6	45	16.2	0.1	3.2	346.2	124.6
	ZAF-6	14–20?	Juvenile	f	Tibia	B		Gold or bronze ring and/or pearl	–19.5	8.2	4.7	6.2	47.4	17.2	0.1	3.2	364.6	132.3
	ZAF-7	18–30	Adult	m	Femur	/			–16.7	8.6	2.4	1.6	42.3	15	0.1	3.3	325.4	115.4
	ZAF-8	25–35	Adult	f	Fibula	C		Gold or bronze ring and/or pearl	–18.4	8.9	4.6	1.6	42.6	15.3	0.1	3.2	304.3	109.3
Valais: Sion Ancienne Placette	ZAF-9	20–40	Adult	f	Ulna	C		Gold or bronze ring and/or pearl	–17.5	9.0	3.8	3.0	48	17.4	0.1	3.2	342.9	124.3
	ZAF-11	25–35	Adult	m	Tibia	C		Weapon	–16.9	7.9	6.8	5.6	39	16.7	0.1	2.7	354.5	151.8
	ZAF-12	>50	Senile?	m	Femur	C	396–211 BC		–18.0	8.6	6.1	5.0	43.6	16.4	0.1	3.1	396.4	149.1
	ZAF-13	nd	nd	nd	Femur	B-C			–19.3	8.2	5.3	1.6	45.4	16.5	0.1	3.2	324.3	117.9
	ZAF-15	nd	nd	f	Humerus	B-C (C)	392–210 BC	Gold or bronze ring and/or pearl	–17.9	9.0	3.8	6.0	46.4	16.9	0.1	3.2	356.9	130.0
	ZAF-19	25+	Adult	f	Femur	/		Gold or bronze ring and/or pearl	–17.7	8.8	3.3	0.6	37.9	13.1	0.1	3.4	315.8	109.2
	ZAF-21	25+	Adult	m	Femur	C			–17.8	9.6	4.6	1.6	48.3	17.4	0.1	3.2	345.0	124.3
	ZAF-22	17–25	Adult	m	Skull	/			–19.4	8.1	3.1	25.4	45.3	15.9	0.2	3.3	283.1	99.4
	ZAF-24	50+	Senile	f	Femur	B/C			–18.6	9.2	8.7	52.8	47.8	17.4	0.1	3.2	341.4	124.3
	ZAF-27	25–35	Adult	m	Femur	D			–18.6	9.5	3.4	1.6	44.9	16	0.2	3.3	299.3	106.7
	ZAF-28	25+	Adult	m	Fibula	C			–18.0	9.5	8.3	6.2	43	16.8	0.1	3.0	358.3	140.0
	SA4	20–40	Adult	f	Skull	HA (C2)	914–811 BC		–16.3	9.0	2.3	5.8	45.0	17.0	0.2	3.1	300.0	113.3
	SA5	nd	nd	m	Long bone (radius-ulna)	/			–15.7	5.3	0.9	5.4	26.1	13.0	0.1	2.3	326.3	162.5
Valais: Sion Nouvelle Placette	SA6	nd	nd	nd	Skull	HA	914–814 BC	Weapon	–16.8	8.5	0.4	2.7	43.0	15.8	0.1	3.2	358.3	131.7
	SN1	20–49	Adult	m	Skull	D1			–18.8	8.4	–0.5	3.0	44.6	17.5	0.1	3.0	371.7	145.8
	SN2	40+	Mature	m	Skull	D1			–17.9	10.0	4.4	1.0	23.2	7.8	0.1	3.5	193.3	65.0
Valais: Sion Parking Remparts	SN3	20–49	Adult	m	Skull	C2			–17.7	10.1	6.1	2.5	32.6	14.7	0.1	2.6	407.5	183.8
	SN4	30–59	Mature	m	Long bone	C2			–18.3	8.6	1.7	2.0	40.8	15.3	0.1	3.1	370.9	139.1
	SN5	30–59	Mature	f	Skull	D2			–19.6	9.7	3.7	1.8	43.6	15.7	0.1	3.2	396.4	142.7
	SP1	30–59	Mature	m	Femur	B-C	391–207 BC		–16.9	8.5	2.5	2.6	46.2	17.0	0.1	3.2	420.0	154.5
	SP11	30–59	Mature	nd	Skull	C1			–18.2	9.4	1.8	2.8	44.9	17.0	0.1	3.1	449.0	170.0
	SP12	20–29	Adult	m	Skull	B-C	384–205 BC		–18.9	6.8	5.5	7.9	46.2	17.0	0.1	3.2	420.0	154.5
	SP13	20–39	Adult	f	Skull	B-C			–18.2	9.3	1.8	2.7	43.8	16.6	0.1	3.1	336.9	127.7

Table 2 (continued)

Burial sites	Number	Age class	Age	Sex	Sample	Archaeological dating	¹⁴ C dating (2)	Grave goods	δ ¹³ C [‰]V-PDB	δ ¹⁵ N [‰]AIR	δ ³⁴ S [‰]V-CDT	%coll	%C	%N	%S	C/ N mol	C/S	N/S
Valais: Sion Passage de la Matze	SP14	5–9	Infant II	nd	Skull	D1	394–209 BC		–19.0	9.8	5.4	4.9	41.6	14.9	0.1	3.3	462.2	165.6
	SP15	30–59	Mature	nd	Fibula?	/			–17.7	9.9	5.4	2.6	45.4	16.7	0.1	3.2	567.5	208.8
	SP16	30–59	Mature	nd	Femur	/			–18.9	9.7	6.6	2.5	35.8	11.6	0.1	3.6	275.4	89.2
	SP17	30–59	Mature	nd	Skull	C1			–18.0	9.6	2.5	3.3	43.2	16.9	0.1	3.0	540.0	211.3
	SP18	40+	Mature	m	Skull	C1			–18.2	8.0	–0.1	1.8	43.1	15.9	0.1	3.2	538.8	198.8
	SP19	60+	Senile	m	Skull	C2			–18.1	9.4	4.8	1.8	40.5	15.0	0.1	3.1	578.6	214.3
	SP27	40–60	Mature	m	Skull	C1			–17.9	9.5	5.9	3.1	44.9	16.6	0.1	3.2	449.0	166.0
	SM1	18–30	Adult	m	Skull	/			–18.8	8.1	4.1	8.3	38.9	17.1	0.1	2.7	324.2	142.5
	SM2	nd	nd	nd	Skull	/			–19.0	9.6	0.5	4.5	39.8	14.8	0.1	3.1	284.3	105.7
	SPC2	5–9	Infant II	nd	Skull	B1		400–212 BC	–19.5	8.6	2.0	1.2	37.3	14.7	0.1	3.0	373.0	147.0
Valais: Sion Petit-Chasseur	SPC4	30–59	Mature	f	Skull	Augustus	349–54 BC	Coins	–20.4	8.8	4.2	2.0	31.8	9.2	0.1	4.0	353.3	102.2
	SPC5	20–39	Adult	f	Scapula	D1		Weapon	–19.1	9.9	2.4	2.1	38.1	15.8	0.1	2.8	317.5	131.7
	SPC6	20–29	Adult	m	Skull	C–D			–18.5	9.1	2.5	2.8	48.2	17.7	0.1	3.2	482.0	177.0
	SPC7	50–60	Mature	f	Skull	D1			–19.0	10.0	3.8	2.4	34	12.4	0.1	3.2	425.0	155.0
	S297	20–49	Adult	m	Skull	C–D			–18.7	10.1	3.8	2.2	43.3	16	0.1	3.2	433.0	159.0
	S422	20–49	Adult	m	Skull	B–D			–18.1	9.2	2.2	3.6	43.6	16	0.1	3.1	363.3	135.0
Valais: Sion-sous-le-Seex	S424	5–9	Infant II	nd	Skull	C2	353–61 BC	Gold or bronze ring and/or pearl	–19.0	9.0	2.6	5.3	34.3	12.5	0.1	3.2	343.0	125.0
	S426	30–59	Mature	f	Long bone	C2		Gold or bronze ring and/or pearl	–18.5	8.9	2.3	4.7	39.7	15.5	0.1	3.0	397.0	155.0
	S427	5–9	Infant II	nd	Skull	C2		pearl	–18.2	9.4	3.8	5.9	41.2	15.2	0.1	3.2	316.9	116.9
	S429	20–29	Adult	f	Skull	C2		Gold or bronze ring and/or pearl	–18.9	9.2	2.3	4.2	44.6	16.6	0.1	3.1	405.5	150.9
	S430	40+	Mature	f	Skull	D1			–19.0	7.9	2.2	4.5	36.7	15.3	0.1	2.8	333.6	139.1
	S431	60+	Senile	f	Skull	/			–19.3	9.8	4.7	1.2	40.0	14.5	0.1	3.2	285.7	103.6
	S432	20–29	Adult	f	Mandibula	C2			–18.9	9.6	4.1	4.8	43.7	16.2	0.1	3.1	397.3	147.3
	S436	30–59	Mature	nd	Humerus?	D1			–18.2	9.5	3.3	3.0	44.1	16.8	0.1	3.1	400.9	152.7
	S437	30–59	Mature	m	Femur	C1			–18.2	8.2	3.8	6.5	45.3	16.6	0.1	3.2	348.5	127.7
	S438	20–39	Adult	f	Fibula	D1			–20.0	7.3	4.9	4.0	44.5	17.9	0.1	2.9	404.5	162.7
	S439	30–59	Mature	m	Fibula	D1	390–156.4	Weapon	–19.0	9.2	3.6	3.0	45.6	16.6	0.1	3.2	380.0	138.3
	S528	20–39	Adult	f	Long bone (radius/ulna)	D1		Gold or bronze ring and/or pearl	–18.5	10.1	5.5	3.7	42.9	17.2	0.1	2.9	390.0	156.4
	S533	30–59	Mature	nd	Skull	D1			–18.5	8.9	–2.0	5.5	42.2	16.0	0.1	3.1	383.6	145.5
	S535	30–59	Mature	nd	Scapula	/			–18.3	8.1	–1.0	4.1	42.5	16.9	0.2	2.9	265.6	105.6
	S537	20–29	Adult	f	Long bone (radius/ulna)	/			–16.3	10.7	4.8	7.1	45.2	16.9	0.1	3.1	376.7	140.8
	S538	30–59	Mature	f	Skull	C2		Coins	–18.2	8.6	1.8	3.9	43.2	16.5	0.1	3.1	308.6	117.9
	S539	20–39	Adult	f	Femur	C2			–18.7	9.3	1.7	5.8	46.0	16.8	0.1	3.2	460.0	168.0
	S540	40+	Mature	f	Skull	C1			–18.4	9.8	2.3	5.0	35.1	12.7	0.1	3.2	438.8	158.8
	S542	20–39	Adult	m	Humerus	C2			–18.4	9.0	3.0	7.4	45.3	16.6	0.1	3.2	566.3	207.5
	S546	>30	Adult	m	Skull	D1		Coins and gold or bronze ring and/or pearl	–19.2	9.1	2.6	4.6	40.1	15.6	0.1	3.0	572.9	222.9
	S547	<50	Adult	f	Skull	C2			–18.8	9.5	3.2	5.0	35.9	13.2	0.1	3.2	398.9	146.7

Table 2 (continued)

Burial sites	Number	Age	Age class	Sex	Sample	Archaeological dating	¹⁴ C dating (2)	Grave goods	δ ¹³ C [‰]v-PDB	δ ¹⁵ N [‰]AIR	δ ³⁴ S [‰]v-CDT	%coll	%C	%N	%S	C/ N mol	C/S	N/S
	S548a	>50	Mature	m	Fibula	D1		Weapon	-18.7	9.9	3.7	2.0	43.4	16.2	0.1	3.1	542.5	202.5
	S548b	<50	Adult	m	Fibula	D1			-18.7	8.3	2.6	3.9	46.8	17.0	0.1	3.2	520.0	188.9
	S549	20–39	Adult	f	Scapula	D1		Gold or bronze ring and/or pearl	-18.4	9.5	1.0	2.9	45.0	16.6	0.1	3.2	500.0	184.4
	S551	60+	Senile	m	Fibula	B-D	360–125 BC		-18.2	9.3	0.3	1.6	41.8	15.5	0.1	3.1	418.0	155.0
Valais: Bramois Panoë	BP-1	30–59	Mature	m	Skull	/			-18.6	10.4	1.8	3.7	37.3	13.5	0.1	3.2	310.8	112.5
	BP-2	40–60	Mature	m	Skull	/			-17.1	9.7	5.9	3.2	37.6	14.3	0.1	3.1	289.2	110.0
	BP3	40+	Mature	f	Skull	B2			-18.2	8.5	3.0	2.4	44.0	16.8	0.1	3.1	338.5	128.5
	BP4	30–59	Mature	f	Skull	D1			-16.7	9.6	6.5	4.3	44.9	16.5	0.1	3.2	345.4	126.9
	BP5	20–49	Adult	m	Skull	D1		Weapon	-18.6	10.5	4.9	3.0	41.8	16.0	0.1	3.0	348.3	133.3
	BP6	20–49	Adult	f	Skull	D2			-19.1	10.2	5.5	1.4	36.6	15.3	0.1	2.8	305.0	127.5
	BP7	50–60	Mature	m	Skull	C-D	191–47 BC		-17.9	10.6	5.1	1.7	42.4	16.4	0.1	3.0	326.2	126.2
	BP8	50–60	Mature	f	Skull	D1		Coins	-18.4	9.1	5.3	2.2	44.4	16.0	0.1	3.2	341.5	123.1
	BP9	60+	Senile	m	Skull	/			-19.0	7.1	4.6	1.3	34.5	14.9	0.1	2.7	313.6	135.5
	BP10	60+	Senile	m	Skull	C1			-18.2	9.5	5.1	4.3	42.2	15.1	0.1	3.3	376.7	137.3
	BP12	40+	Mature	m	Femur	B-C (C1)	348–55 BC		-18.1	8.1	2.7	2.3	41.6	16.5	0.1	2.9	341.5	137.5
	BP13	60+	Senile	f	Long bone (radius/ulna)	D2			-18.0	9.9	5.6	4.4	43.4	16.4	0.1	3.1	313.6	149.1
	BP14	20–29	Adult	nd	Skull	B-C	389–208 BC		-17.9	10.5	7.6	5.3	41.7	15.9	0.1	3.1	379.1	144.5
Valais: Bramois Villa Lathion-Lopes	BV28	60+	Senile	m	Skull	/			-18.5	9.9	4.2	2.2	49.4	17.9	0.1	3.2	380.0	137.7
Valais: Bramois Villa Schaller	BV25	20–49	Adult	m	Fibula	/			-17.3	9.1	7.4	3.7	43.0	16.4	0.1	3.1	358.3	136.7
	BV26	30–59	Mature	m	Skull	C-D	190–41 BC		-18.3	9.9	7.1	2.1	42.7	16.5	0.2	3.0	284.7	110.0
	BV27	60+	Senile	m	Skull	/			-18.2	10.1	5.4	2.4	43.7	17.9	0.1	2.8	437.0	179.0
Grisons: Trun - Darvella	TDA-20	30–40	Adult	m	Skull	B/C			-19.7	8.8	4.0	2.1	42.6	15.8	0.2	3.1	284.0	105.3
	TDA-21	40–60	Mature	m	Skull	B/C			-19.4	8.4	4.3	5.5	44.2	16.1	0.1	3.2	315.7	115.0
	TDA-22	20–35	Adult	f	Skull	B/C			-18.9	8.1	4.5	5.1	42.0	15.4	0.1	3.2	323.1	118.5
Grisons: Castaneda	CAS-6	<18	Juvenil	nd	Skull	A-B (C?)	513–393		-16.3	8.3	6.2	2.9	42.1	15.5	0.1	3.2	323.8	119.2
Bern: Engelhalbinsel (Retschenbachstrasse)	15	Animal			Pig	C-D			-21.4	4.8	8.0	9.7	47.4	17.4	0.1	3.9	395.0	145.0
	26				Ovicaprid	C-D			-20.4	6.6	7.7	3.0	43.5	16.3	0.1	3.6	334.6	125.4
	7 (1)				Ovicaprid	C-D	193–55		-21.2	4.3	6.3	2.6	44.6	16.3	0.1	3.7	371.7	135.8
	7 (2)				Cattle	B-D	348–61		-21.5	6.0	4.8	4.0	47.0	17.4	0.1	3.9	391.7	145.0
Grisons: Bonaduz	4	Animal			Ovicaprid	A-C	535–111		-21.3	2.3	-0.3	2.4	42.6	15.7	0.1	3.2	355.0	130.8

Selected grave goods are shown

m male; f female; nd not determined

sites of the Swiss Plateau, the area of Bern in particular, noticeably more female than male burials were found (Fig. 2). This, however, seems to be a “Bernese phenomenon” which was previously described by Jud and Ulrich-Bochsler (2014). Biased excavation techniques and collection practices at the beginning of the twentieth century could have caused the unbalanced sex distribution with a loss of skeletal material and a lack of information on burial findings from that time. Different mortuary practices for males and females might have also led to the specific sex distribution as found in Engehalbinsel. Iron Age human remains were also found in Ipsach (Ramstein 2010; Zweifel 2015), and burials were discovered in Belp and Niederwichttrach (Schoch and Ulrich-Bochsler 1987; Suter and Ulrich-Bochsler 1984). Geographical clusters within the findings’ distribution suggest two centers around the Belp Mountain including Belp, Münsingen, Niederwichttrach and around the Enge peninsula. Different burial sites in the area of Bümpliz, a suburb of Bern, revealed Late Iron Age burials (Schoch and Ulrich-Bochsler 1987; Stähli 1977). Additionally, burials in Deisswil, a district of the Bernese suburb Stettlen, were excavated (Rey 1999). In total, samples of 74 individuals were taken from burial sites within the region of Bern for stable isotope analysis (Table 1). Previous published stable isotope data from the burial site of Münsingen (Moghaddam et al. 2016) are part of the analyses for comparison.

The village of Andelfingen is situated in the north of the Canton of Zurich, between Winterthur and Schaffhausen (Fig. 1). It was believed that the Late Iron Age burial site here might have been connected to a later Celtic oppidum. Nine male and 12 female graves were


found (Viollier 1912). A total of 16 samples could be collected (Table 1).

Iron Age burial sites in the Swiss Alps

The Swiss Alps in the south of the country stretches from Geneva in the west to the Austrian border in the east. Almost 60% of the Swiss landscape is covered by the Alps. During the last century, many Iron Age burials were found in the Swiss Alps including the burial sites of Sion, Bramois, Castaneda and Trun Darvella (Kaenel 1999). Sion and Bramois lie between the Pennine and the Bernese Alps (Fig. 1), where a great number of Iron Age burials were found but destroyed in the nineteenth century before documentation. Burials were excavated in Sion, Sous le Scex Passage de la Matze, at the center of Sion at Nouvelle Placette and in the Avenue du Petit Chasseur in Sion (Curdy et al. 2009; Debard 2014). Furthermore, 3 km from Sous le Scex further burials were found at Bramois (Fig. 1) and excavated during recent years (Curdy et al. 2009). Findings of animal bones lead to the assumption that a settlement was located nearby. In total, samples of 70 individuals were taken from within the region of Valais for the stable isotope analysis (Table 1).

The area of Grisons is entirely mountainous. Several burials were excavated in the Darvella village at the Rhine headwaters (Fig. 1). The burial site might have belonged to a larger settlement (Tanner 1980). The village of Castaneda is located at an ancient transalpine route that connected the Raetia with the southern Alpine valleys via the Little St. Bernhard. Four samples from Grisons could be used for this study (Keller-Tarnuzzer 1933; Nagy 2008).

Fig. 2 Distribution of sex and age of all individuals grouped in the Swiss Plateau and Swiss Alps



Methods

Morphologic–anthropological analysis

The age at death of adults was established through analyzing the closure of the cranial sutures (Szilvássy 1988), age-related changes of the pubic symphysis (Acsádi and Nemeskéri 1970) and also the dental surface wear (Brothwell 1981). The stages of the epiphyseal union (Ferembach et al. 1979), long bone measurements (Scheuer and Black 2000; Schmid and Künle 1958) and eruption of deciduous and permanent teeth were studied for non-adults (Ubelaker 1989). The individuals were assigned to the following age classes:

Infants Ia (0–3 years); Infants Ib (4–7 years); Infants II (8–14 years); Juvenile (15–18/20 years); Adult (20–45 years); Mature (46–60 years); Senile (61 years and over).

For sex estimation, features on the Os coxae including the ventral arc, subpubic concavity, medial surface of the ischiopubic ramus and differences in the greater sciatic notch were analyzed (Ferembach et al. 1979; Grupe et al. 2015; Herrmann et al. 1990; Sjøvold 1988). Cranial morphology such as the nuchal crest, mastoid process, supra-orbital margin and the mental eminence were observed (Buikstra and Ubelaker 1994) and femoral head diameters were measured, after Herrmann et al. (1990). Sexual dimorphic features are not fully developed in subadults and therefore no sex estimation was performed.

For some sites morphological analyses were verified and/or additional information was taken from publications (Debard 2014; Hug 1956; Schoch and Ulrich-Bochsler 1987; Ulrich-Bochsler and Rüttimann 2014; Viollier 1912).

The faunal remains have been published by Rehazek and Nussbaumer (2014).

Stable isotope analysis

Most of the samples were collected from crania to have similar bone turnover rates. The collagen extraction of all samples ($n = 164$) was conducted by modified methods of Ambrose (1993) and Longin (1971). For detailed information of the extraction method and stable isotope analysis, see [supplement material](#).

Collagen with less than 1% in proportion to the dry weight and samples with a molar C/N relation beyond a 2.9–3.6 range were excluded. Furthermore, %C, %N and %S values that strongly diverge from recent collagen values (C: 43%, N: 15–16% and S: 0.2%) were not taken into consideration (Ambrose 1990, 1993; DeNiro 1985; Nehlich and Richards 2009).

Dating

Most burial sites were dated through archaeological studies (Debard 2014; Jud and Ulrich-Bochsler 2014; Rey 1999; Stähli 1977; Suter and Ulrich-Bochsler 1984; Viollier 1912).

Additionally, selected samples were radiocarbon-dated after Szidat et al. (2016) (Department of Archaeometrie of the Curt-Engelhorn-Center in Mannheim, Germany; Department of Chemistry and Biochemistry at the University of Bern, Switzerland).

Statistics

The individuals were divided into two geographical groups, the Swiss Plateau and Swiss Alpine regions. Furthermore, statistical analyses were carried out for groups of sex and age.

The results were analyzed via SPSS Statistics 23. Tests of normality were performed through Kolmogorov–Smirnov. Non-parametric tests (Mann–Whitney- U test and Kruskal–Wallis test) for independent samples were used due to the sample size and not normal distribution of the data (Fagerland 2012). As non-parametric tests are less robust than t tests, a rejection of the null hypothesis is more assured, while the same outcome can be observed in parametric tests. Nevertheless, analysis of variance (t test and ANOVA) for independent groups was performed to verify the non-parametric tests. Multivariate testing was conducted through the $\delta^{13}\text{C}/\delta^{15}\text{N}$ -ratio in order to analyze differences in the distribution of the values.

Null hypothesis stating that the mean stable isotope values of categorical groups are equal was rejected with $p < 0.05$. The level of significance is indicated as following: $p \leq 0.05$ (*); $p \leq 0.01$ (**); $p \leq 0.001$ (***)

Results

Morphological and individual data

A total of 52 adult individuals were aligned as male, 66 as female and 19 non-subadults as indeterminate. Age estimation for subadults revealed 5 infant Ia, 13 infants Ib, and 9 infants II. In total, 90 individuals were sampled from the Swiss Plateau and 74 individuals from the Swiss Alps (Fig. 2). Results of radiocarbon dating of selected samples were assigned to dating phases (Table 2).

Descriptive statistics of total stable isotope data

Due to collagen quality criteria, 129 out of 164 samples could be further analyzed for stable carbon, nitrogen and sulfur isotopes. Descriptive statistics of all stable isotope data are shown in Table 3.

In total, $\delta^{13}\text{C}$ values of all individuals range from -21.5‰ to -16.1‰ with a mean of -18.7‰ and a standard deviation (SD) of 1.0‰ . The $\delta^{15}\text{N}$ values range from 6.8‰ to 11.3‰ (mean = 8.8‰ ; SD = 0.9‰). The $\delta^{34}\text{S}$ range from -2.0‰ to 8.7‰ (mean = 4.2‰ ; SD = 2.2‰).

Table 3 Descriptive statistics of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$ with means, standard deviation (*SD*), median, minimum (*Min*) and maximum (*Max*)

Geography	Region	Site	n	δ ¹³ C [‰] _{V-PDB}				δ ¹⁵ N [‰] _{AIR}				δ ³⁴ S [‰] _{V-CDT}						
				Mean	SD	Median	Min	Max	Mean	SD	Median	Min	Max	Mean	SD	Median	Min	Max
Münsingen			63	-19.5	0.7	-19.6	-20.8	-17.4	8.8	0.5	8.8	7.6	10.2	1.0	2.2	0.4	-2.3	7.6
Swiss Plateau	Bern	Engelhalbinsel	25	-18.0	1.2	-18.0	-20.0	-16.3	8.2	1.1	8.0	6.9	11.3	5.7	1.2	5.8	1.6	7.1
		Ipsach	4	-17.9	1.2	-18.0	-19.2	-16.3	9.8	0.8	9.9	8.9	10.7	5.7	1.4	6.2	3.6	6.7
		Belp	7	-18.7	1.3	-18.9	-20	-16.1	8.4	0.5	8.3	7.8	9.3	2.9	1.7	3.3	0.2	5.1
		Niederwichttrach	3	-20.4	0.9	-20.1	-21.5	-19.7	8.8	0.02	8.8	8.8	8.8	2.3	0.3	2.3	2	2.5
		Bümpliz	11	-18.5	0.9	-18.2	-20.3	-17.6	7.7	0.5	7.6	7.0	8.4	6.1	1.4	6.5	2.1	7.1
		Stettlen-Deisswil	8	-19.3	0.6	-19.3	-19.9	-18.6	8.9	0.7	8.7	8.2	10.4	2.3	0.9	2.1	1.3	3.8
		Total	58	-19.1	0.9	-19.2	-21.5	-16.1	8.4	1.0	8.2	6.9	11.3	4.8	2.0	5.5	0.2	7.1
	Andelfingen		14	-18.2	0.8	-18.1	-19.5	-16.7	8.9	0.5	8.9	8.1	9.6	5.4	2.2	4.6	2.4	8.7
Total	72	-18.9	1.0	-19.1	-21.5	-16.1	8.5	0.9	8.4	6.9	11.3	4.9	2.0	5.4	0.2	8.7		
Swiss Alps	Valais	Sion	41	-18.4	0.8	-18.4	-20.0	-16.3	9.0	0.8	9.2	6.8	10.7	2.7	1.9	2.5	-2.0	5.9
		Bramois	12	-18.0	5.3	-18.1	-18.6	-16.7	9.6	0.8	9.7	8.1	10.6	5.4	1.6	5.2	2.7	7.6
		Total	53	-18.3	0.7	-18.3	-20.0	-16.3	9.2	0.8	9.3	6.8	10.7	3.3	2.2	3.3	-2.0	7.6
	Grisons	Castaneda	1	-16.3	/	8.3	/	6.2	/									
		Trun Darvella	3	-19.3	0.4	-19.4	-19.7	-18.9	8.4	0.4	8.4	8.1	8.8	4.3	2.5	4.3	4.0	4.5
		Total	4	-18.6	1.6	-19.2	-19.7	-16.3	8.4	0.3	8.4	8.1	8.8	4.8	1.0	4.4	4.0	6.2
Total	57	-18.3	0.8	-18.3	-20.0	-16.3	9.1	0.8	9.2	6.8	10.7	3.4	2.1	3.7	-2.0	7.6		
Overall total	129	-18.7	1.0	-18.7	-21.5	-16.1	8.8	0.9	8.8	6.8	11.3	4.2	2.2	4.6	-2.0	8.7		
sex and age	Male	42	-18.5	0.8	-18.5	-20.0	-16.7	8.8	0.9	8.9	6.8	10.6	3.9	2.1	3.8	-5.0	8.5	
	Female	54	-18.6	0.9	-18.8	-20.0	-16.1	8.6	0.9	8.6	6.9	10.7	4.5	2.1	4.7	1.0	8.7	
	Infants II	5	-19.4	0.8	-19.5	-20.2	-18.2	8.5	1.2	8.6	7.0	9.8	4.5	1.6	5.4	2.0	5.8	
	Infants Ib	10	-19.4	0.7	-19.5	-20.3	-18.1	8.9	1.0	8.6	7.6	10.7	5.6	1.3	6.2	2.1	6.3	
	Infants Ia	3	-18.0	1.5	-18.7	-19.1	-16.3	10.6	0.6	10.3	10.2	11.3	5.6	1.8	6.6	3.6	6.7	
	nd	15	-18.4	1.3	-18.2	-21.5	-16.3	8.9	0.9	8.8	7.2	10.5	3.0	2.8	2.5	-2.0	7.6	

Data for Münsingen are presented in a separate row for comparison. All data are divided into geography, region, and sites. Totals are shown separately for geography and regions. Results of sex and age are shown with the overall total in bold

Stable isotope data for the faunal remains range from -21.5 to -20.2‰ for $\delta^{13}\text{C}$ (mean = -21.1‰ ; SD = 0.4‰), 2.3‰ to 6.6‰ (mean = 4.8‰ ; SD = 1.5‰) for $\delta^{15}\text{N}$ and -0.3‰ to 8.0‰ (mean = 5.3‰ ; SD = 3.0‰) for $\delta^{34}\text{S}$ (Table 2).

Statistical tests of stable isotope data

Results of the statistical tests are shown in Table 4. The geographical sites of the Swiss Plateau and the Swiss Alps showed highly significant differences considering all stable isotope data ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$ $p = 0.000***$). A significant higher $\delta^{13}\text{C}$ mean for the Swiss Alps was observed (Table 3). In total, 27 individuals showed enriched $\delta^{13}\text{C}$ values (higher than or equal to -18.0‰) 11 were female, 9 male, 4 non-determinable adults, 1 infant Ia and 1 juvenile (Fig. 3); 15 derive from the Swiss Plateau, and 12 from the Swiss Alps (Table 2). Highly significant differences were also observed for the $\delta^{15}\text{N}$ values with the highest mean for the Swiss Alpine regions (Fig. 4). Regarding the distribution of the $\delta^{13}\text{C}/\delta^{15}\text{N}$ -ratio, a high significance ($p = 0.000***$) was also obtained.

Analyses of the different sexes revealed no significance regarding the two different geographical areas (Table 4). The evaluation of each burial site also separately revealed no differences. Additionally, $\delta^{13}\text{C}/\delta^{15}\text{N}$ -ratio for sex was not significant for the Swiss Alps ($p = 0.315$) and the Swiss Plateau ($p = 0.452$), whereas on adding the Münsingen individuals to the Swiss Plateau significant distinct mean values could be observed within the Swiss Plateau ($\delta^{13}\text{C}$, $p = 0.011*$; $\delta^{15}\text{N}$, $p = 0.006**$). Additionally, differences between the sexes from all burial sites in the $\delta^{15}\text{N}$ values ($p = 0.006**$) with males having higher $\delta^{15}\text{N}$ values (mean = 9.0‰) than females (mean = 8.6‰) could be observed.

Non-parametric tests showed differences in the $\delta^{13}\text{C}$ ($p = 0.042*$), $\delta^{15}\text{N}$ ($p = 0.003**$) and $\delta^{34}\text{S}$ ($p = 0.006**$) mean values between the different age classes of the whole dataset (excluding Münsingen), but individuals from the Swiss Alpine regions showed no differences between the age classes. In contrast, the individuals within the Swiss Plateau showed high significances in the mean values ($\delta^{13}\text{C}$ $p = 0.045*$; $\delta^{15}\text{N}$ $p = 0.002**$; $\delta^{34}\text{S}$ $p = 0.018*$).

Discussion

Consumption of millet in Late Iron Age Switzerland

The data of the human remains indicate an overall diet mainly based on animal protein and C_3 plant sources. The data are in agreement with previously published isotopic data from Iron Age sites of present-day Germany (Knipper et al. 2014; Oelze et al. 2012). In total, the $\delta^{13}\text{C}$ data show a relatively wide range, both in the Swiss Plateau ($\Delta 5.0\text{‰}$) and in the Swiss

Alpine habitat ($\Delta 3.7\text{‰}$), and some individuals show a clear enrichment in ^{13}C . This variation shows that there must have been a big diversity in plant food resources in these societies, especially in C_3 and C_4 plants. The reason for this could have been, e.g., different food preferences and also different food distribution. Various migration patterns in these societies might also be a reason for the observed variations. Values greater than -18‰ indicate a significant intake of C_4 -plants in the diet, such as millet (Le Huray and Schutkowski 2005). In this study, 21% of the individuals show enriched $\delta^{13}\text{C}$ values, both in the Swiss Plateau (15/72) and in the Swiss Alps (12/57). The frequency of enriched specimens is highly related to the site: e.g., Zürich Andelfingen (7/14) and Bern Bümpliz (5/11) have frequencies of 50 and 45%, respectively (both Swiss Plateau). These frequencies imply a regular consumption of millet in these populations compared to, e.g., Bern Engehalbinsel with none of the individuals revealing values higher than or equal to -18.0‰ . Different frequencies of enriched individuals are also shown for the Valais (Swiss Alps), e.g., for Sion 17% (7/41) and 33% (4/12) for Bramois. However, the statistics indicate an overall higher intake of C_4 plant for populations from the Swiss Alpine regions. As no animal samples from these areas could be analyzed, and the faunal–human trophic level cannot be reconstructed, it remains unclear whether this result might reflect a direct consumption of millet and/or a consumption of animal proteins originating from animals fed on C_4 plants. For the Iron Age, isotopic evidence for millet consumption in continental Europe has been shown by, e.g., Knipper et al. (2014), Le Huray and Schutkowski (2005), and Lightfoot et al. (2012). Evidence of, especially, broomcorn millet (*Panicum miliaceum*) in Late Iron Age Switzerland was mentioned by Jacomet and Jacquat (1999). Broomcorn millet as one of the earliest crops was first cultivated in Asia and reached Eastern Europe in the first millennium BC (Miller et al. 2016). Hence, the cultivation of millet in Late Iron Age Switzerland must have been affected through influences from other regions, especially the Mediterranean. Other studies confirm that millet has been cultivated in, e.g., Late Bronze Age Greece (Petroutsas and Manolis 2010) and in northern Greek Early Iron Age sites (Papathanasiou et al. 2013). Furthermore, it is known that there were intense economic exchanges from Massalia to surrounding areas (Bouby et al. 2011). From findings in Mediterranean France, it has been reported that Greek ceramics and wine amphorae also played a major role in imports (Loughton 2009) as well as agricultural resources. Among different rivers, the Rhône has been a main trade route from the Mediterranean to east-central Gaul spreading Greek cultural influences. Hence, different kinds of seeds were probably brought from the Mediterranean world to the Celtic culture.

Climatic variability might also have had an effect on farming as warmer climates favor the growth of C_4 plants

Table 4 Statistical non-parametric Mann–Whitney *U* and Kruskal–Wallis tests for independent groups

	Isotope	Swiss Plateau vs. Swiss Alps	Male vs. female	Male vs. female Swiss Plateau	Male vs. female Swiss Alps	Age classes	Age classes Swiss Plateau	Age classes Swiss Alps	Swiss Plateau vs. Swiss Alps (males)	Swiss Plateau vs. Swiss Alps (females)
Non-parametric <i>p</i> value	$\delta^{13}\text{C}$	0.000***	0.308	0.984	0.793	0.042*	0.045*	0.109	0.282	0.101
	$\delta^{15}\text{N}$	0.000***	0.115	0.234	0.417	0.003**	0.002**	0.272	0.042*	0.000***
	$\delta^{34}\text{S}$	0.000***	0.181	0.416	0.914	0.006*	0.018*	0.481	0.306	0.022*

Significance: * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$

(Ehleringer et al. 1991). These variabilities could have been due to regional climatic differences (Van Klinken et al. 2000), but also due to climate changes through time. The climate of Switzerland which is also influenced by the Atlantic Ocean shows great variability among different regions. The west-east-oriented Alps lead to relatively strong gradients in the climate (Wanner et al. 1997) and act as a barrier between the moderate European and the Mediterranean climate. Southern Switzerland, influenced by Mediterranean weather, is therefore characterized by a much milder climate than the north (Brönnimann et al. 2014). C_4 plants such as millet are more drought-tolerant than C_3 plants and therefore its cultivation is easier in warmer climates including with short growing seasons (Lightfoot et al. 2015). In fact, this could have favored the cultivation of C_4 plants in southern areas of Switzerland. A connection between millet cultivation and climate variation was also observed by Jacob et al. (2009). Their study indicated

climate deterioration with a decline of agricultural work during the Bronze Age–Iron Age transition in the French Prealps.

For Münsingen, the $\delta^{13}\text{C}$ values were more positive in the later compared to the earlier phases (Moghaddam et al. 2016). Including Münsingen in the dataset, the highest mean (-18.6‰) was observed in individuals dating to LT C (Kruskal–Wallis $p = 0.000***$). The highest $\delta^{13}\text{C}$ value overall was found in an adult female from Bern Belp (BEB-A2272) dating to the LT C period. A climate change during the Late Iron Age with a warmer climate in the Late La Tène phases should be taken into consideration which could have also facilitated the cultivation of C_4 plants. However, isotopic variability between plants and within plants can be very large even over short distances (Van Klinken et al. 2000). Non-photosynthetic tissues of plants such as roots are generally ^{13}C -enriched (Cernusak et al. 2009), which might have played a minor role in the diet for some populations.




Fig. 3 Stable carbon and nitrogen isotope data, means and standard deviation (SD) of different sexes and subadults. Additionally, means and SD of Münsingen males and females are shown




Fig. 4 Stable carbon and nitrogen isotope data, means and standard deviation (SD) of the different sites. Additionally, mean and SD of Münsingen, Swiss Plateau, and Swiss Alps are shown

The question emerges whether millet might have been part of a “low” or “high” status food. The data do not support any correlation between social status expressed by precious grave goods and millet consumption. Some burials contained no grave goods at all, others large amounts of jewelry (IP133, ZAF5 and ZAF 9). No difference between the sexes could be observed, and individuals with a signal of C_4 plant consumption were of both sexes and all ages (Fig. 3).

The evaluation of the age classes revealed significant differences for the Swiss Plateau. A mean of -18.7‰ for the females shows a ^{13}C -enrichment of almost 1‰ in the infants Ia (mean -18.0‰) probably indicating a breastfeeding signal (Fuller et al. 2006).

The intake of animal protein and social status

The overall $\delta^{15}\text{N}$ data show a relatively wide range, both in the Swiss Plateau ($\Delta 4.4\text{‰}$) including all individuals from the Bern region and in the Swiss Alpine habitat ($\Delta 3.9\text{‰}$). This variation indicate a high diversity in food resources in these societies, in particular in the distribution of animal proteins. The nitrogen isotope values of the animals show that the omnivore pig lies within the range of the herbivores (Table 2). It is likely that domestic pigs were fed similarly to herbivores, as found for Roman Age Italy (Prowse et al. 2004). The $\delta^{15}\text{N}$ mean of the humans is about 4.4‰ higher than that of the herbivores and agrees with a difference of one trophic level

(Hedges and Reynard 2007; Schwarcz and Schoeninger 1991). Differences were observed between the Swiss Plateau and Swiss Alps where the individuals showed significantly higher $\delta^{15}\text{N}$ values (Fig. 4). When assuming a general shift of 3‰ between trophic levels (Schwarcz and Schoeninger 1991), the 5‰ difference in the herbivores for the burial sites in the region of Valais (Sion and Bramois) correspond to a shift of almost two levels. This indicates a diet higher in animal proteins of these populations in comparison to the others. It could be evidence for consumption of dairy products and/or meat in these Alpine regions. This corresponds to the recently published study by Carrer et al. (2016), even if their data are not directly comparable to this study. They identified dairy lipids on Iron Age ceramics from the high Alps, indicating the earliest evidence of dairy production in the Alps.

The intake of freshwater fish for the Valais populations should be mentioned as the sites are close to the River Rhône. Nevertheless, a significant consumption of freshwater fish is less likely as the $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ values reflect a more terrestrial-based diet.

Depending on soil types, climate and land use, the $\delta^{15}\text{N}$ can vary widely (Goude and Fontugne 2016; Van Klinken et al. 2000). The “manuring effect”, however, should be taken into account. Manuring is known to have been practiced since prehistoric times (Bakels 1997; Nielsen and Kristiansen 2014). The populations in the southern Alpine areas compared to the Swiss Plateau probably had distinct patterns of crop

cultivation and husbandry influenced by different geological conditions and climatic variabilities.

The Münsingen population revealed sex-specific dietary differences with males having a diet higher in animal protein (Moghaddam et al. 2016). None of the other sites revealed a similar result. The Münsingen population indeed had a distinct social structure based on sex. However, the question remains whether the higher $\delta^{15}\text{N}$ values of the Swiss Alpine populations might derive from a larger number of male samples (Fig. 2), as males had generally higher $\delta^{15}\text{N}$ (Fig. 3), but the analyses of the females exclusively still showed significant differences ($p = 0.000***$) in the mean $\delta^{15}\text{N}$ (9.3‰). Therefore, the higher values of the Alpine regions do not derive from a larger number of males.

There was no correlation between grave goods, such as weapons and jewelry, with $\delta^{15}\text{N}$ values, in contrast to Münsingen and Iron Age Bohemia (Le Huray and Schutkowski 2005). However, a higher mean for burials with weapons ($n = 6$; $\delta^{15}\text{N}$, mean = 9.0‰) was observed compared to other adult and older individuals ($n = 101$, $\delta^{15}\text{N}$ mean = 8.7‰; Table 2).

The means of the infants Ia and adult females result in a shift of almost one trophic level ($\Delta^{15}\text{N} = 2\text{‰}$). According to previous published studies, this result agrees with a breastfeeding signal (Beaumont et al. 2015; Katzenberg 2008; Katzenberg et al. 1996)

Migration in the Late Iron Age

Sulfur data of the ovicaprids showed a relatively wide range. The pig showed the highest value of 8.0‰ (Table 2). Since there is just little fractionation between consumer and food (Jay et al. 2013; Nehlich et al. 2011), the human sulfur values indicate a terrestrial-based diet for Iron Age Switzerland. It should be kept in mind that the faunal data represent more or less the local value. This hampers the interpretation between the faunal and human samples from different regions.

Comparing the sites, there are highly significant differences (Fig. 5) as $\delta^{34}\text{S}$ values reflect local geological data and indicate different local $\delta^{34}\text{S}$ values (Vika 2009). In total, the $\delta^{34}\text{S}$ data show the highest variations, both in the Swiss Plateau ($\Delta 8.5\text{‰}$) and in the Swiss Alpine regions ($\Delta 10.7\text{‰}$). Comparing sites from the Bern, Zurich and Grisons regions, no significant differences were observable (Kruskal–Wallis; $p = 0.832$). The populations from Valais differ significantly with lower $\delta^{34}\text{S}$ values compared to other sites. The significantly different local signature becomes more evident by comparing the data within the area of Grisons (Table 1). The samples from Trun Darvella show little variation while the sample from Castaneda is higher. The significantly different value of the ovicaprids from Bonaduz (near Trun Darvella) shows local differences in the area of Grisons. The sulfur standard deviation shows higher variations in all sites compared to their $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values.




Fig. 5 Stable carbon and sulfur isotope data, means and standard deviation (SD) of the different sites. Additionally, mean and SD of Münsingen, Swiss Plateau, and Swiss Alps are shown

The reasons for this are different small-scale geological formations of these regions but it could also show migration of single individuals within the societies. Significant differences could be observed between the sites within the Valais with Sion having lower $\delta^{34}\text{S}$ than Bramois (Table 3). The high SD for the area of Valais most likely derives from local specific sulfur isotope differences. The same could be observed for the individuals in the area of Bern. The $\delta^{34}\text{S}$ of the different sites in Bern, with Engthalbinsel, Ipsach and Bümpliz, have more positive means than Stettlen-Deisswil and Niederwichttrach (Table 3), which is very likely due to regional–geographical variabilities (Fig. 1). Nevertheless, mobility in Iron Age Switzerland is obvious, as has already been published for Münsingen (Moghaddam et al. 2013; Scheeres et al. 2016). In the case of Andelfingen, the data differ significantly in their $\delta^{34}\text{S}$ values (Fig. 5). Their variances might derive through geological distinction of the soil which is reflected by the plants grown at that place. A certain proportion of freshwater fish in the diet is proposed for individuals with higher values than 8.0‰, the maximum value for terrestrial animals. The high amount of methionine in fish protein also reveals a higher amount of methionine in the consumers' amino acid compound. This causes a reflection of higher values in the consumer even with a small intake of fish (Nehlich et al. 2010).

Limitation of the study

The sex and age determination was analyzed by morphological features of the bones. Even though the maximum number of features were analyzed, in some cases the skeletons were not complete. This could have led to uncertainties, especially for ageing older individuals, and thus might have had a minor effect on the statistical analysis. For more valid data, morphological analyses were carried out in comparison with previous published analyses and archaeological data. In cases of high uncertainties, the individuals were assigned as not determined.

Additionally, more animal bones would be needed to create a convenient food-web. Faunal remains from the Swiss Alpine regions, in particular, would be necessary to obtain information about the intake of animal protein.

Conclusion

This study suggests great differences in the subsistence strategies of Swiss Late Iron Age populations from different regions. Geological conditions with different environments, climate and cultural aspects probably led to differences in agricultural practice and animal husbandry. Isotopic analyses indicate dietary differences between geographical regions that were also caused by migration during the Iron Age. The exchange with the Mediterranean world probably had a great influence on the Iron Age culture of this region. Sex-related

dietary differences were found only in Münsingen which led to the assumption that this population may have differed from the populations studied here in their socio-economic structure.

The presented isotopic data provide a vast overview of the Swiss Late Iron Age and are relevant for future studies of prehistoric populations. These isotopic data are indispensable for projects on the spread of millet in prehistoric central Europe.

Acknowledgements We would like to thank the archaeological services of the Cantons of Bern, Valais, and Grisons, and the Anthropological Institute at the University of Zurich for support. We appreciate the cooperation of Geneviève Perréard, Julie Debar, Marco Millela, Thomas Reitmaier, Viera Trancik, and Albert Hafner. The authors also thank Inga Siebke and Christine Cooper for assistance and Matthias Dörmötör for image editing. We are very grateful to both anonymous reviewers for their helpful comments and advices.

Compliance with ethical standards

Grant sponsorship This research was supported by the Swiss National Science foundation (CR13I3_149583).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Acsádi G, Nemeskéri J (1970) History of human lifespan and mortality. Akadémiai Kiadó, Budapest
- Ambrose SH (1990) Preparation and characterization of bone and tooth collagen for isotopic analysis. *J Archaeol Sci* 17:431–451
- Ambrose SH (1993) Isotopic analysis of paleodiets: methodological and interpretive considerations. In: Sanford MK (ed) *Investigations of ancient human tissue - chemical analyses in anthropology*. Gordon and Breach, Amsterdam, pp 59–130
- Ambrose SH, Norr L (1993) Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: Lambert JB, Grupe G (eds) *Prehistoric human bone: archaeology at the molecular level*. Springer, Berlin, pp 1–37
- Bakels CC (1997) The beginnings of manuring in western Europe. *Antiquity* 71:442–445
- Beaumont J, Montgomery J, Buckberry J, Jay M (2015) Infant mortality and isotopic complexity: New approaches to stress, maternal health, and weaning. *Am J Phys Anthropol* 157:441–457. doi:10.1002/ajpa.22736
- Bocherens H, Drucker D (2003) Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems. *Int J Osteoarchaeol* 13:46–53
- Bollongino R et al (2013) 2000 years of parallel societies in stone age central Europe. *Science* 342:479–481. doi:10.1126/science.1245049
- Bouby L, Boissinot P, Marinval P (2011) Never mind the bottle. Archaeobotanical evidence of beer-brewing in Mediterranean France and the consumption of alcoholic beverages during the 5th century BC. *Hum Ecol* 39:351–360

- Brönnimann S et al (2014) Climate change in Switzerland: a review of physical, institutional, and political aspects. *Clim Change* 5:461–481. doi:[10.1002/wcc.280](https://doi.org/10.1002/wcc.280)
- Brothwell DR (1981) Digging up bones; the excavation, treatment and study of human skeletal remains, third edn. Oxford University Press, Oxford
- Buikstra JE, Ubelaker DH (1994) Standards for data collection from human skeletal remains: proceedings of a seminar at the Field Museum of Natural History, organised by Jonathan Haas. vol 44. Arkansas Archeological Survey, Fayetteville, Arkansas
- Carrer F et al. (2016) Chemical Analysis of Pottery Demonstrates Prehistoric Origin for High-Altitude Alpine Dairying. *PLoS ONE* 11:ARTN e0151442 doi: [10.1371/journal.pone.0151442](https://doi.org/10.1371/journal.pone.0151442)
- Cernusak LA et al (2009) Viewpoint: Why are non-photosynthetic tissues generally C-13 enriched compared with leaves in C-3 plants? Review and synthesis of current hypotheses. *Funct Plant Biol* 36: 199–213. doi:[10.1071/Fp08216](https://doi.org/10.1071/Fp08216)
- Collis J (1984) The European iron age. Butler & Tanner, London
- Craig OE, Ross R, Andersen SH, Milner N, Bailey GN (2006) Focus: sulphur isotope variation in archaeological marine fauna from northern. *Eur J Archaeol Sci* 33:1642–1646
- Curdy P, Mariéthoz F, Pernet L, Rast-Eicher A (2009) Rituels funéraires chez les Séduens. Les nécropoles du second âge de fer en valais central (IV-I. siècle av. J.-C.). In: Curdy P (ed) Cahiers D'Archéologie Romande: Archaeologica Vallesiana 3, vol 112. Cahiers D'Archéologie Romande Lausanne, p 304
- Debard J (2014) Approche bioanthropologique des conditions socio-économiques à la fin de l'âge du Fer en Suisse occidentale : stature, croissance et stress environnemental., Université de Genève
- DeNiro MJ (1985) Postmortem preservation and alteration of in vivo bone-collagen isotope ratios in relation to paleodietary reconstruction. *Nature* 317:806–809
- Dupras TL, Schwarcz HP (2001) Strangers in a strange land: stable isotope evidence for human migration in the Dakhleh oasis. *Egypt J Archaeol Sci* 28:1199–1208
- Ehleringer JR, Sage RF, Flanagan LB, Pearcy RW (1991) Climate change and the evolution of C4 photosynthesis. *Trends Ecol Evol* 6:95–99. doi:[10.1016/0169-5347\(91\)90183-X](https://doi.org/10.1016/0169-5347(91)90183-X)
- Fagerland MW (2012) t-tests, non-parametric tests, and large studies-a paradox of statistical practice? *Bmc Med Res Methodol* 12 doi:Artn 78 Doi [10.1186/1471-2288-12-78](https://doi.org/10.1186/1471-2288-12-78)
- Ferembach D, Schwidetzky I, Stloukal M (1979) Recommendations for age and sex diagnoses of skeletons. *Bull Mém Soc Anthropol Paris* 6:7–45
- Fornander E, Eriksson G, Lidén K (2008) Wild at heart: approaching pitted ware identity, economy and cosmology through stable isotopes in skeletal material from the neolithic site Korsnas in Eastern Central Sweden. *J Anthropol Archaeol* 27:281–297
- Fuller BT, Molleson TI, Harris DA, Gilmour LT, Hedges REM (2006) Isotopic evidence for breastfeeding and possible adult dietary differences from Late/Sub-Roman Britain. *Am J Phys Anthropol* 129:45–54
- Fuller BT, De Cupere B, Marinova E, Van Neer W, Waelkens M, Richards MP (2012) Isotopic reconstruction of human diet and animal husbandry practices during the Classical-Hellenistic, imperial, and Byzantine periods at Sagalassos. *Turk Am J Phys Anthropol* 149:157–171
- Goude G, Fontugne M (2016) Carbon and nitrogen isotopic variability in bone collagen during the Neolithic period: Influence of environmental factors and diet. *J Archaeol Sci* 70:117–131. doi:[10.1016/j.jas.2016.04.019](https://doi.org/10.1016/j.jas.2016.04.019)
- Grupe G, Harbeck M, McGlynn GC (2015) Prähistorische anthropologie. Springer, Berlin
- Hedges REM, Reynard LM (2007) Nitrogen isotopes and the trophic level of humans in archaeology. *J Archaeol Sci* 34:1240–1251
- Herrmann B, Grupe G, Hummel S, Piepenbrink H, Schutkowski H (1990) Prähistorische Anthropologie. Leitfaden der Feld- und Labormethoden [Prehistoric Anthropology. Guide of field and laboratory methods]. Springer, Berlin
- Hoefs J (2009) Stable isotope geochemistry, 6th edn. Springer, Berlin
- Howcroft R, Eriksson G, Lidén K (2012) Conformity in diversity? Isotopic investigations of infant feeding practices in two iron age populations from Southern Oland, Sweden. *Am J Phys Anthropol* 149:217–230. doi:[10.1002/ajpa.22113](https://doi.org/10.1002/ajpa.22113)
- Hug E (1956) Die Anthropologische Sammlung im Naturhistorischen Museum Bern, vol 13. Naturhistorisches Museum Bern, Bern
- Jacob J, Disnar JR, Arnaud F, Gauthier E, Billaud Y, Chapron E, Bardoux G (2009) Impacts of new agricultural practices on soil erosion during the Bronze Age in the French Prealps Holocene 19:241–249 doi:[10.1177/0959683608100568](https://doi.org/10.1177/0959683608100568)
- Jacomet S, Jacquat C (1999) Ackerbau: Bedeutung der Anbaupflanzen und ihre mögliche Verwendung. In: Müller F, Kaenel G., Lüscher, G. (ed) Die Schweiz vom Paläolithikum bis zum frühen Mittelalter. SPM IV. Eisenzeit. Verlag Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel, pp 105–109
- Jay M, Richards MP (2006) Diet in the iron age cemetery population at Wetwang Slack, East Yorkshire, UK: carbon and nitrogen stable isotope evidence. *J Archaeol Sci* 33:653–662
- Jay M, Montgomery J, Nehlich O, Towers J, Evans J (2013) British Iron Age chariot burials of the Arras culture: a multi-isotope approach to investigating mobility levels and subsistence practices. *World Archaeol* 45:473–491. doi:[10.1080/00438243.2013.820647](https://doi.org/10.1080/00438243.2013.820647)
- Jud P, Ulrich-Bochsler S (2014) Bern, Reichenbachstrasse. Die Gräber aus dem Oppidum auf der Engehalbinsel Bern. Archäologischer Dienst des Kantons Bern, Bern
- Kaenel G (1999) Die Archäologie der Eisenzeit in der Schweiz. In: Müller F, Kaenel G, Lüscher G (eds) Die Schweiz vom Paläolithikum bis zum frühen Mittelalter. SPM IV. Eisenzeit. Verlag Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel, pp 15–23
- Katzenberg MA (2008) Stable isotope analysis: a tool for studying past diet, demography, and life history. In: Katzenberg MA, Saunders SR (eds) Biological anthropology of the human skeleton. Wiley-Liss, New Jersey, pp 413–441
- Katzenberg MA, Herring DA, Saunders SR (1996) Weaning and Infant Mortality: Evaluating the Skeletal Evidence vol 39. Yearbook of Physical Anthropology 39:177–199
- Katzenberg MA, McKenzie HG, Losey RJ, Gorunova OI, Weber A (2012) Prehistoric dietary adaptations among hunter-fisher-gatherers from the Little Sea of Lake Baikal, Siberia, Russian Federation. *J Archaeol Sci* 39:2612–2626. doi:[10.1016/j.jas.2011.08.010](https://doi.org/10.1016/j.jas.2011.08.010)
- Keller-Tarnuzzer K (1933) Die eisenzeitliche Siedlung von Castaneda: Grabung 1932 Anzeiger für schweizerische Altertumskunde: Neue Folge. Indicateur d'antiquités Suisse: Nouv Sér 35:161–177
- Kinaston RL, Buckley HR, Gray A (2013) Diet and social status on Taumako, a Polynesian outlier in the Southeastern Solomon Islands. *Am J Phys Anthropol* 151:589–603
- Knipper C et al (2014) Social differentiation and land use at an Early Iron Age “princely seat”: bioarchaeological investigations at the Glauberg (Germany). *J Archaeol Sci* 41:818–835
- Knipper C et al. (2016) What is on the menu in a Celtic town? Iron Age diet reconstructed at Basel-Gasfabrik, Switzerland. *J Archaeol Anthropol Sci*:1–20 doi:[10.1007/s12520-016-0362-8](https://doi.org/10.1007/s12520-016-0362-8)
- Koon HEC, Nicholls RA (2016) Investigating heterogeneity in diet across early Iron Age assemblages from Slovenia using stable isotopes. *Am J Phys Anthropol* 159:195–196
- Le Huray JD, Schutkowski H (2005) Diet and social status during the Late period in bohemia: carbon and nitrogen stable isotope analysis of bone collagen from kutna hora-karlovo and radovesice. *J Anthropol Archaeol* 24:135–147
- Le Huray JD, Schutkowski H, Richards DA (2006) La tène dietary variation in central Europe: a stable isotope study of human skeletal

- remains from bohemia. In: Gowland R, Knüsel C (eds) *Social archaeology of funerary remains*. Oxbow, Oxford, pp 99–122
- Lee-Thorp JA (2008) On isotopes and Old bones. *Archaeometry* 50:925–950. doi:10.1111/j.1475-4754.2008.00441.x
- Lightfoot E, Šlaus M, O'Connell TC (2012) Changing cultures, changing cuisines: cultural transitions and dietary changing in iron Age, roman and early Mediaeval Croatia. *Am J Phys Anthropol* 148:543–556
- Lightfoot E, Šlaus M, Šikanjić PR, O'Connell TC (2015) Metals and millets: bronze and iron Age diet in inland and coastal Croatia seen through stable isotope analysis. *Archaeol Anthropol Sci* 7:375–386. doi:10.1007/s12520-014-0194-3
- Longin R (1971) New method of collagen extraction for radiocarbon dating. *Nature* 230:241–242
- Lösch S, Grupe G, Peters J (2006) Stable isotopes and dietary adaptations in humans and animals at Pre-Pottery Neolithic Nevali Cori, south-east Anatolia. *Am J Phys Anthropol* 131:181–193
- Loughton ME (2009) Getting smashed: the deposition of amphorae and the drinking of wine in Gaul during the late iron Age. *Oxf J Archaeol* 28:77–110. doi:10.1111/j.1468-0092.2008.00320.x
- Miller NF, Spengler RN, Frachetti M (2016) Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene* 26:1566–1575. doi:10.1177/0959683616641742
- Moghaddam N, Langer R, Ross S, Nielsen E, Lösch S (2013) Multiple osteosclerotic lesions in an Iron Age skull from Switzerland (320–250 BC) - an unusual case. *Swiss Med Wkly* 143:ARTN w13819 doi: 10.4414/smw.2013.13819
- Moghaddam N, Müller F, Hafner A, Lösch S (2016) Social stratigraphy in Late Iron Age Switzerland: Stable carbon, nitrogen and sulphur isotope analysis of human remains from Münsingen. *Archaeol Anthropol Sci* 8:149–160. doi:10.1007/s12520-014-0221-4
- Müller F (1996) Latènezeitliche Grabkeramik aus dem Berner Aaretal. In: Niffeler U (ed) *Jahrbuch der Schweizerischen Gesellschaft für Ur- und Frühgeschichte*, vol 79. Schweizerischen Gesellschaft für Ur- und Frühgeschichte, Basel, pp 43–66
- Müller F, Lüscher G (2004) *Die Kelten in der Schweiz*. Konrad Theiss, Stuttgart
- Müller F, Kaenel G, Lüscher G (1999) *Die Schweiz vom Paläolithikum bis zum frühen Mittelalter*. SPM IV. Eisenzeit. Verlag Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel
- Nagy P (2008) *Das eisenzeitliche Gräberfeld von Castaneda*, Kanton Graubünden. University of Zurich
- Naumann E, Price TD, Richards MP (2014) Changes in dietary practices and social organization during the pivotal late iron age period in Norway (AD 550–1030): isotope analyses of Merovingian and Viking Age human remains. *Am J Phys Anthropol* 155:322–331. doi:10.1002/ajpa.22551
- Nehlich O, Richards MP (2009) Establishing collagen quality criteria for sulphur isotope analysis of archaeological bone collagen. *Archaeol Anthropol Sci* 1:59–75
- Nehlich O, Borić D, Stefanović S, Richards MP (2010) Sulphur isotope evidence for freshwater fish consumption: a case study from the Danube Gorges, SE Europe. *J Archaeol Sci* 37:1131–1139. doi:10.1016/j.jas.2009.12.013
- Nehlich O, Fuller BT, Jay M, Mora A, Nicholson RA, Smith CI, Richards MP (2011) Applications of sulphur isotope ratios to examine weaning patterns and freshwater fish consumption in Roman Oxfordshire, UK. *Geochim Cosmochim Acta* 75:4963–4977
- Nehlich O, Fuller BT, Márquez-Grant N, Richards MP (2012) Investigation of diachronic dietary patterns on the islands of Ibiza and formentera, Spain: evidence from sulfur stable isotope ratio analysis. *Am J Phys Anthropol* 149:115–124
- Nielsen NH, Kristiansen SM (2014) Identifying ancient manuring: traditional phosphate vs. multi-element analysis of archaeological soil. *J Archaeol Sci* 42:390–398. doi:10.1016/j.jas.2013.11.013
- Oelze VM et al (2012) Multi-isotopic analysis reveals individual mobility and diet at the Early Iron Age monumental tumulus of Magdalenenberg, Germany. *Am J Phys Anthropol* 148:406–421. doi:10.1002/ajpa.22063
- Papathanasiou A, Panagiotopoulou E, Beltsios K, Papakonstantinou MF, Sipsi M (2013) Inferences from the human skeletal material of the Early Iron Age cemetery at Agios Dimitrios, Fthiotis, Central Greece. *J Archaeol Sci* 40:2924–2933
- Petroutsas EI, Manolis SK (2010) Reconstructing Late Bronze Age diet in mainland Greece using stable isotope analysis. *J Archaeol Sci* 37: 614–620. doi:10.1016/j.jas.2009.10.026
- Privat KL, O'Connell TC, Hedges REM (2007) The distinction between freshwater- and terrestrial-based diets: methodological concerns and archaeological applications of sulphur stable isotope analysis. *J Archaeol Sci* 34:1197–1204
- Prowse T, Schwarcz HP, Saunders S, Macchiarelli R, Bondioli L (2004) Isotopic paleodiet studies of skeletons from the imperial roman-age cemetery of Isola Sacra, Rome, Italy. *J Archaeol Sci* 31:259–272
- Ramstein M (2010) Ipsach, Räberein - Latènezeitliche Gräber im römischen Gutshofareal. In: *ArchBE. Jahrbuch des Archäologischen Dienstes des Kantons Bern, Fundberichte 2009*. Erziehungsdirektion des Kantons Bern, Bern, pp 96–97
- Redfern RC, Hamlin C, Athfield NB (2010) Temporal changes in diet: a stable isotope analysis of late Iron Age and Roman Dorset, Britain. *J Archaeol Sci* 37:1149–1160
- Redfern RC, Millard AR, Hamlin C (2012) A regional investigation of subadult dietary patterns and health in late Iron Age and Roman Dorset, England. *J Archaeol Sci* 39:1249–1259
- Rehazek A, Nussbaumer M (2014) Tierknochen. In: Jud P, Ulrich-Bochsler S (eds) *Bern, Reichenbachstrasse - Neue Gräber aus dem Latènezeitlichen Oppidum auf der Engehalbinsel*. Archaeologischer Dienst Bern, Rub Media, Bern, pp 47–49
- Rey T (1999) Das latènezeitliche Gräberfeld von Stettlen-Deisswil BE. In: Niffeler U (ed) *Jahrbuch der Schweizerischen Gesellschaft für Ur- und Frühgeschichte*, vol 82. Schweizerischen Gesellschaft für Ur- und Frühgeschichte, Basel, pp 117–148
- Richards MP, Hedges REM, Molleson TI, Vogel JC (1998) Stable isotope analysis reveals variations in human diet at the Poundbury Camp cemetery site. *J Archaeol Sci* 25:1247–1252. doi:10.1006/j.jasc.1998.0307
- Richards MP, Fuller BT, Hedges REM (2001) Sulphur isotopic variation in ancient bone collagen from Europe: implications for human palaeodiet, residence mobility, and modern pollutant studies. *Earth Planet Sci Lett* 191:185–190. doi:10.1016/S0012-821x(01)00427-7
- Richards MP, Fuller BT, Sponheimer M, Robinson T, Ayliffe L (2003) Sulphur isotopes in palaeodietary studies: a review and results from a controlled feeding experiment. *Int J Osteoarchaeol* 13:37–45
- Scheeres M, Knipper C, Schönfelder M, Hauschild M, Siebel W, Alt KW (2016) Bioarchaeometric investigations (87Sr/86Sr AND $\delta^{18}O$) of the La tène burial community of münsingen-rain; Switzerland. In: Arnold B (ed) *The oxford handbook of the archaeology of the continental Celts*. Oxford University Press, Oxford
- Scheuer L, Black S (2000) *Developmental juvenile osteology*. Academic, London
- Schmid F, Künle F (1958) Das Längenwachstum der langen Röhrenknochen in bezug auf Körperlänge und Lebensalter Fortschritte auf dem Gebiet der Röntgenstrahlen und der Nuklearmedizin 89:350–356
- Schoch W, Ulrich-Bochsler S (1987) *Die Anthropologische Sammlung des Naturhistorischen Museums Bern - Katalog der Neueingänge 1956–1985*. Jahrb Naturhist Mus Bern 9:267–350
- Schoeninger MJ, DeNiro MJ (1984) Nitrogen and carbon isotopic composition of bone-collagen from marine and terrestrial animals. *Geochim Cosmochim Acta* 48:625–639. doi:10.1016/0016-7037(84)90091-7

- Szwarcz HP, Schoeninger MJ (1991) Stable isotope analyses in human nutritional ecology. *Yearb Phys Anthropol* 34:283–321
- Sjøvold T (1988) Geschlechtsdiagnose am Skelett In: Knussmann R (ed) *Anthropologie: Handbuch der vergleichenden Biologie des Menschen* Gustav Fischer Verlag, Stuttgart, New York, pp 444–480
- Stähli B (1977) *Schriften des Seminars für Urgeschichte der Universität Bern*, vol 3
- Stevens RE, Hedges REM (2004) Carbon and nitrogen stable isotope analysis of northwest European horse bone and tooth collagen, 40,000 BP-present: palaeoclimatic interpretations. *Quat Sci Rev* 23: 977–991. doi:[10.1016/j.quascirev.2003.06.024](https://doi.org/10.1016/j.quascirev.2003.06.024)
- Stevens RE, Jacobi RM, Higham TFG (2010) Reassessing the diet of upper palaeolithic humans from Gough's cave and Sun hole, cheddar gorge, Somerset, UK. *J Archaeol Sci* 37:52–61
- Stöckli WE (1995) Das spätlatènezeitliche Gäberfeld von Niederwichtach-Bachtelen im Kanton Bern. In: Schmid-Sikimić B, and Della Casa P, editors. *Tran Europam Beiträge zur Bronze- und Eisenzeit zwischen Atlantik und Altai Festschrift für Margarita Primas*. Bonn: Dr. Rudolf Habelt GmbH, p 251–264
- Suter P, Ulrich-Bochsler S (1984) Neuere Mittellatène-Grabkomplexe aus dem Kanton Bern. Ein Beitrag zur Latène C-Chronologie des schweizerischen Mittellandes. In: Osterwalder C (ed) *Jahrbuch Schweizerische Gesellschaft für Ur- und Frühgeschichte*, vol 67. Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel, pp 73–93
- Szidat S, Vogel E, Gubler R, Lösch S (2016) Radiocarbon dating of bones at the LARA laboratory at Bern *Radiocarbon*:12 doi:[10.1017/RDC.2016.90](https://doi.org/10.1017/RDC.2016.90)
- Szilvássy J (1988) Altersdiagnose am Skelett In: Knussmann R (ed) *Anthropologie: Handbuch der vergleichenden Biologie des Menschen*, Gustav Fischer Stuttgart, pp 421–435
- Tanner A (1980) *Das Latènegräberfeld von Trun-Darvella*. Schriften des Seminars für Urgeschichte der Universität Bern, vol 5. Historisch-Archäologischer, Zürich
- Ubelaker DH (1989) *Human skeletal remains: excavation, analysis, interpretation*, 2nd edn. Taraxacum, Washington D.C
- Ulrich-Bochsler S, Rüttimann D (2014) Die anthropologie. In: Jud P, Ulrich-Bochsler S (eds) *Bern, Reichenbachstrasse - Neue Gräber aus dem Latènezeitlichen Oppidum auf der Engehalbinsel*. Archaeologischer Dienst Bern, Rub Media, Bern, pp 68–90
- Van Klinken GJ, Richard MP, Hedges REM (2000) An overview of causes for stable isotopic variations in past European human populations: environmental, ecophysiological, and cultural effects. In: Ambrose SH, Katzenberg MA (eds) *Biogeochemical approaches to paleodietary analysis*. Kluwer, New York, pp 39–63
- Vika E (2009) Strangers in the grave? Investigating local provenance in a Greek Bronze Age mass burial using delta S-34 analysis. *J Archaeol Sci* 36:2024–2028
- Viollier D (1912) Fouille exécutées par les soins du Musée National. VII. Le cimetière gallo-helvète d'Andelfingen (Zurich) *Indicateur d'antiquités suisses: Nouvelle série* 14:16–57. doi:[10.5169/seals-158989](https://doi.org/10.5169/seals-158989)
- Wanner H, Rickli R, Salvisberg E, Schmutz C, Schüepp M (1997) Global climate change and variability and its influence on Alpine climate - Concepts and observations. *Theor Appl Climatol* 58:221–243. doi:[10.1007/Bf00865022](https://doi.org/10.1007/Bf00865022)
- Zweifel U (2015) Fünf Latènezeitliche Gräber aus Ipsach, Räberein. Mit einem Beitrag zu latènezeitlichen Kinderbestattungen im Schweizer Mittelland. In: Bern ADK (ed) *Jahrbuch des Archäologischen Dienstes des Kantons Bern* 2015. Rub Media, Bern, pp 112–135